

**Healthfirst: Health Home 2023 GIV Measure Details**  
**Updated 9/5/23**

CMA Action	Abbreviation	Measure Name	Replacement Measure	Abbreviation	Measure Description	Additional Notes
Set-up Dental Appointment	ADV	Annual Dental Visit	Oral Evaluation, Dental Services	OED	The percentage of members 2-20 years of age who had at least one dental visit during the measurement year.	
Set-up PCP visit for Medication Adherence	AMM - Acute Phase	Antidepressant Medication Mgmt - Acute Phase			The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment for at least 84 days.	
Set-up PCP visit for Medication Adherence	AMM - Continuation Phase	Antidepressant Medication Mgmt - Continuation Phase			The percentage of members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an antidepressant medication treatment for at least 180 days.	
Set-up PCP visit for Medication Adherence	AMR	Asthma Medication Ratio			The percentage of members ages 5-64 who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.	Replaces the Medication Management for Asthma measures
Set-up PCP visit for Medication Monitoring	APM	Metabolic Monitoring for Children and Adolescents on Antipsychotics			The percentage of children and adolescents 1-17 years of age who had two or more antipsychotic prescriptions and had metabolic testing.	
Set-up visit for Medical Screening	BCS	Breast Cancer Screening			The percentage of women 50-74 years of age who had a mammogram to screen for breast cancer.	
Set-up visit for Medical Screening	CCS	Cervical Cancer Screening			The percentage of women 21-64 years of age who were screened for cervical cancer using either of the following criteria: <ul style="list-style-type: none"> <li>Women 21-64 who had cervical cytology performed every 3 years</li> <li>Women 30-64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years</li> </ul>	
Set-up PCP visit for Diabetic Monitoring	CDC	Comprehensive Diabetes Care Blood Pressure Control (<140/90 mm Hg)	Blood Pressure Control for Patients with Diabetes	BPD	The members most recent blood pressure reading taken during the measurement year is less than 140/90 mm Hg.	
Set-up PCP visit for Diabetic Monitoring	CDC	Comprehensive Diabetes Care HbA1c Poor Control (>9.0%)	Hemoglobin A1c Control with Diabetes (>9.0%)	HBD	The members most recent HbA1c level is greater than 9.0% or is missing a result, or if an HbA1c test was not done during the measurement year.	
Set-up PCP visit for Diabetic Monitoring	CDC	Comprehensive Diabetes Care HbA1c Control (<8.0%)	Hemoglobin A1c Control with Diabetes (<8.0%)	HBD	The members most recent HbA1c level is less than 8.0%	
Set-up PCP visit for Diabetic Monitoring	CDC	Comprehensive Diabetes Care Hemoglobin A1c (HbA1c) Testing	Hemoglobin A1c Control with Diabetes	HBD	An HbA1c test performed during the measurement year	
Set-up Eye Exam	CDC	Comprehensive Diabetes Care: Eye Exam (retinal) Performed	Eye Exam for Patients with Diabetes	EED	Screening or monitoring for diabetic retinal disease as identified by administrative data. This includes diabetics who had one of the following: <ul style="list-style-type: none"> <li>A retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) in the measurement year</li> <li>A negative retinal or dilated eye exam (negative for retinopathy) by an eye care professional in the year prior to the measurement year.</li> </ul>	
Set-up PCP visit for Diabetic Monitoring	CDC	Comprehensive Diabetes Care: Medical Attention for Nephropathy	Kidney Health Evaluation for Patients Diabetes	KED	A nephropathy screening or monitoring test or evidence of nephropathy, as documented through administrative data. This includes diabetics who had one of the following during the measurement year: <ul style="list-style-type: none"> <li>A nephropathy screening or monitoring test</li> <li>Evidence of treatment for nephropathy or ACE/ARB therapy</li> <li>Evidence of stage 4 chronic kidney disease</li> <li>Evidence of end stage renal disease</li> <li>Evidence of kidney transplant</li> <li>A visit with a nephrologist</li> <li>At least one ACE inhibitor or ARB dispensing event</li> </ul>	
Set-up visit for Medical Screening	CHL	Chlamydia Screening in Women			The percentage of women 16-24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.	
Set-up visit for Medical Screening	COL	Colorectal Cancer Screening			The percentage of members 50-75 years of age who had appropriate screening for colorectal cancer.	
Set-up SUD Provider Visit and Confirm Attendance	IET	Initiation and Engagement of Alcohol and Other Drug Abuse or Dependence Treatment			Initiation of AOD Tx: The percentage of members ages 13+ who initiate treatment for AOD within 14 days of diagnosis.  Engagement of AOD Tx: The percentage of members ages 13+ who initiated treatment and who were engaged in ongoing AOD treatment within 34 days of the initiation visit.	
Set-up visit for Medication Adherence	QPODD	Use of Pharmacotherapy for Alcohol Abuse or Dependence			The percentage of members ages 18+ with an encounter associated with alcohol use or dependence, with at least 1 prescription for appropriate pharmacotherapy at any time during the measurement year.	
Set-up visit for Medication Adherence	QPODN	Initiation of Pharmacotherapy upon New Episode of Opioid Dependence			The percentage of members ages 18+ who initiate pharmacotherapy with at least 1 prescription of visit for opioid treatment medication within 30 days following a new diagnosis of opioid dependence.	
Set-up PCP visit for Medication Adherence	SAA	Adherence to Antipsychotic Meds for People with Schizophrenia			The percentage of members 19-64 years of age during the measurement year with schizophrenia or schizoaffective disorder who were dispensed and remained on an antipsychotic medication for at least 80% of their treatment period.	
Set-up PCP visit for Cardiovascular Disease Monitoring	SMC	Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia			The percentage of members 18-64 years of age with schizophrenia or schizoaffective disorder and cardiovascular disease, who had an LDL-C test during the measurement year.	
Set-up PCP visit for Diabetic Monitoring	SMD	Diabetes Monitoring for People with Diabetes and Schizophrenia			The percentage of members 18-64 years of age with schizophrenia or schizoaffective disorder and diabetes who had both an LDL-C test and an HbA1c test during the measurement year.	
Set-up PCP visit for Medication Adherence	SPC	Statin Therapy for Patients with Cardiovascular Disease - 80 PDC			The percentage of males 21-75 years of age and females 40-75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD), and remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period.	PDC = Proportion of Days Covered
Set-up PCP visit for Medication Adherence	SPC	Statin Therapy for Patients with Cardiovascular Disease - Received Tx			The percentage of males 21-75 years of age and females 40-75 years of age during the measurement year, who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD), and were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year.	
Set-up PCP visit for Diabetic Monitoring	SSD	Diabetes Screening for People with Schizophrenia or Bipolar Disorder Using Antipsychotic Meds			The percentage of members 18-64 years of age with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year	
Set-up PCP Visit	W30	Well-Child Visit First 30 months (6+ visits)			The percentage of members who had the following number of well-child visits with a PCP during the measurement year. 1. First 15 Months: 6+ well-child visits 2. 15 Months - 30 Months: 2+ well-child visits	Replaces the Wel-Child Visit First 15 Months measure
Set-up PCP Visit	WCV	Child and Adolescent Well Care Visits			The percentage of members ages 3-21 who had at least one well-care visit with a PCP or an OB/GYN practitioner during the measurement year.	Replaces Well-Child Visits in the 3rd, 4th, 5th & 6th Years of Life & Adolescent Well Care Visit measures